Amendment to the Claims

In the Claims:

Please amend Claims 1, 8, 49, and 50 as follows:

- 1. (Currently Amended) A method of constructing a blank library of optically distinct reporter labeled carriers, said method comprising the steps of:
- (a) providing a plurality of carriers, at least some two of which are identical, each carrier being configured to support a plurality of optically distinct reporters and a plurality of compounds of interest, none of the plurality of carriers as initially provided having any reporters or compounds of interest attached thereto;
- (b) providing a plurality of reaction vessels, such that at least one reaction vessel is available for each unique member of the blank library to be constructed;
- (c) providing a plurality of optically distinct reporters, such that at least some two of the plurality of optically distinct reporters are different from each other;
- (d) in each reaction vessel, apportioning at least one carrier, such that any reaction vessel that includes more than one carrier will include only identical carriers;
- (e) in each reaction vessel, apportioning at least one optically distinct reporter, such that a unique combination of the at least one carrier and the at least one optically distinct reporter is achieved in each reaction vessel, and such that the optically distinct reporters in each vessel enable each carrier in that vessel to be distinguished from carrier and reporter combinations in other vessels; and
- (f) attaching said at least one optically distinct reporter to said at least one carrier in each reaction vessel, such that each carrier in the same reaction vessel will have an identical set of optically distinct reporters attached to it, each reaction vessel including a set of optically distinct reporter labeled carriers that is uniquely different from the optically distinct reporter labeled carriers of each other reaction vessel, the plurality of reaction vessels thereby defining a blank library of optically distinct reporter labeled carriers, such that compounds of interest can later be attached to the optically distinct reporter labeled carriers of the blank library.
- 2. (Previously Presented) The method of Claim 1, wherein at least one reaction vessel contains a carrier that is optically distinct from others of said plurality of carriers in other reaction vessels, and wherein no reaction vessel contains a mixture of optically distinct carriers.

29

- 3. (Previously Cancelled)
- 4. (Previously Cancelled)
- 5. (Previously Cancelled)
- 6. (Previously Cancelled)
- 7. (Previously Cancelled)
- 8. (Currently Amended) A method of constructing a blank library of optically distinct reporter labeled carriers, said method comprising the steps of:
- (a) providing a plurality of optically distinct carriers, at least some two of which are identical, each carrier being configured to support a plurality of optically distinct reporters and a plurality of compounds of interest, none of the plurality of carriers as initially provided having any reporters or compounds of interest attached thereto;
- (b) providing a plurality of reaction vessels, such that at least one reaction vessel is available for each unique member of the library to be constructed;
- (c) providing a plurality of optically distinct reporters, such that at least some two of the plurality of optically distinct reporters are different from each other;
- (d) in each reaction vessel, apportioning at least one optically distinct carrier and at least one optically distinct reporter, such that:
- (i) any one of the plurality of reaction vessels that includes more than one optically distinct carrier will include only identical optically distinct carriers; and
- (ii) a unique combination of the at least one optically distinct carrier and the at least one optically distinct reporter is achieved in each reaction vessel; and
- (e) attaching said at least one reporter to said at least one carrier in each reaction vessel, such that each optically distinct carrier in the same reaction vessel will have an identical set of optically distinct reporters attached to it, each reaction vessel including a set of optically distinct reporter labeled carriers that is uniquely different from the optically distinct reporter labeled carriers of each other reaction vessel, the plurality of reaction vessels thereby defining a blank library of optically distinct reporter labeled carriers, such that compounds of interest can later be attached to the optically distinct reporter labeled carriers of the blank library.
- 9. (Previously Presented) The method of Claim 8, wherein no reaction vessel contains a mixture of different optically distinct carriers.

33. (Previously Cancelled)

A 4	/D 1	~
44 /	Previousl	y Cancelled)
J7.		Y Canteentur

- 35. (Previously Cancelled)
- 36. (Previously Cancelled)
- 37. (Previously Cancelled)
- 38. (Previously Cancelled)
- 39. (Previously Cancelled)
- 40. (Previously Cancelled)
- 41. (Previously Presented) The method of Claim 2, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on size.
- 42. (Previously Presented) The method of Claim 2, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on shape.
- 43. (Previously Presented) The method of Claim 2, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on color.
- 44. (Previously Presented) The method of Claim 2, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on an intensity of color.
- 45. (Previously Presented) The method of Claim 8, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on size.
- 46. (Previously Presented) The method of Claim 8, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on shape.
- 47. (Previously Presented) The method of Claim 8, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on color.

///

///

- 48. (Previously Presented) The method of Claim 8, wherein at least one reaction vessel includes optically distinct carriers that can be optically distinguished from carriers in other reaction vessels based on an intensity of color.
- 49. (Currently Amended) A method of constructing a library of optically distinct reporter labeled carriers to which no compounds of interest are yet attached, the method comprising the steps of:
- (a) providing a plurality of carriers, at least some two of which are identical, each carrier being configured to support a plurality of optically distinct reporters and a plurality of compounds of interest, none of the plurality of carriers as initially provided having any reporters or compounds of interest attached thereto;
- (b) providing a plurality of reaction vessels, such that at least one reaction vessel is available for each unique member of the library to be constructed;
- (c) providing a plurality of optically distinct reporters, such that at least some two of the plurality of optically distinct reporters are different from each other;
- (d) in each reaction vessel, apportioning at least one carrier, such that any reaction vessels that includes more than one carrier will include only identical carriers;
- (e) in each reaction vessel, apportioning at least one optically distinct reporter, such that a unique combination of the at least one carrier and the at least one optically distinct reporter is achieved in each reaction vessel, and such that the optically distinct reporters in each vessel enable each carrier in that vessel to be distinguished from carrier and reporter combinations in other vessels; and
- (f) attaching the at least one optically distinct reporter to the at least one carrier in each reaction vessel, such that each carrier in the same reaction vessel will have an identical set of optically distinct reporters attached to it, each reaction vessel including a set of optically distinct reporter labeled carriers that is uniquely different from the optically distinct reporter labeled carriers of each other reaction vessel, the plurality of reaction vessels thereby defining a blank library of optically distinct reporter labeled carriers, each member of the library consisting of a carrier to which at least one optically distinct reporter is attached.
- 50. (Currently Amended) A method of constructing a blank library of optically distinct reporter labeled carriers, the method comprising the steps of:

- (a) providing a plurality of optically distinct reporters, such that at least some two of the plurality of optically distinct reporters are different from each other;
- (b) based on the number of different optically distinct reporters provided, determining all possible binary codes that can be produced using the optically distinct reporters provided;
- (c) providing a plurality of reaction vessels, such that each reaction vessel corresponds to a specific binary code;
- (d) providing a plurality of carriers, at least some two of which are identical, each carrier being configured to support a plurality of optically distinct reporters and a plurality of compounds of interest, none of the plurality of carriers as initially provided having any reporters or compounds of interest attached thereto;
- (e) in each reaction vessel, apportioning the optically distinct reporters required to produce the binary code corresponding to the reaction vessel;
- (f) in each reaction vessel, apportioning at least one carrier, such that any reaction vessel that includes more than one carrier will include only identical carriers; and
- (g) in each reaction vessel, attaching the optically distinct reporters defining the binary code for the reaction vessel to each carrier in the reaction vessel, such that each carrier in the same reaction vessel will have an identical set of optically distinct reporters attached to it, each reaction vessel including a set of optically distinct reporter labeled carriers that is uniquely different from the optically distinct reporter labeled carriers of each other reaction vessel, the plurality of reaction vessels thereby defining a blank library of optically distinct reporter labeled carriers, such that compounds of interest can be later attached to the optically distinct reporter labeled carriers of the blank library.
- 51. (Previously Presented) The method of Claim 50, wherein the step of apportioning the optically distinct reporters required to produce the binary code corresponding to the reaction vessel comprises the step of adding each different type of optically distinct reporter to the reaction vessel separately, such that if more than two different types of optically distinct reporters are required, each additional type of optically distinct reporter is not added to the reaction vessel until the previously added types of optically distinct reporters have combined to achieve a cluster.

///

30 || ///

1	
2	
3	l
4	
5	
6	I
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	
25	
26	
27	
28	
29	
30	١

52. (Previously Presented) The method of Claim 50, wherein the step of providing a plurality of optically distinct reporters comprises the step of providing optically distinct reporters whose sizes are smaller than a spatial resolution limit of an imaging system that will be used to analyze the library.